

General Volume-Preserving Mechanical Systems

Bin Zhou*

Institute of High Energy Physics, Chinese Academy of Sciences,
P. O. Box 918-4, Beijing 100039, P. R. China,

Han-Ying Guo[†]

Institute of Theoretical Physics, Chinese Academy of Sciences,
P. O. Box 2735, Beijing 100080, P. R. China,

and Ke Wu[‡]

Department of Mathematics, Capital Normal University,
Beijing 100037, P. R. China

July 18, 2002

Abstract

In this letter, we present the general form of equations that generate a volume-preserving flow on a symplectic manifold $(\mathcal{M}, \omega)$. It is shown that every volume-preserving flow has some 2-forms acting the rôle of the Hamiltonian functions in the Hamiltonian mechanics and the ordinary Hamilton equations are included as a special case with a 2-form $\frac{1}{n-1} H \omega$ where H is the corresponding Hamiltonian.

In classical mechanics, most of the systems can be described by the ordinary Hamilton equations

$$\dot{q}^i = \frac{\partial H}{\partial p_i}, \quad \dot{p}_i = -\frac{\partial H}{\partial q^i}. \quad (1)$$

*Email: zhoub@ihep.ac.cn

[†]hyguo@itp.ac.cn

[‡]wuke@itp.ac.cn

For an autonomous system, namely, a system with the Hamiltonian manifestly independent of the time t , the Hamiltonian H is a function on the phase space $\mathcal{M}$ which is precisely the cotangent bundle of the configuration space. Abstractly, a Hamiltonian system can be defined on a symplectic manifold $(\mathcal{M}, \omega)$ where $\mathcal{M}$ is a $2n$ -dimensional orientable differentiable manifold endowed with a symplectic 2-form ω at every point of $\mathcal{M}$. For the details of Hamiltonian mechanics on a symplectic manifold, see, for example, [1]. Then, according to Darboux's theorem, for each point x of $\mathcal{M}$, there is a coordinate neighborhood $(U; q^1, \dots, q^n, p_1, \dots, p_n)$ of x such that¹

$$\omega|_U = dp_i \wedge dq^i. \quad (2)$$

Thus a Hamiltonian system with the Hamiltonian H can still be described locally by eqs. (1). The solutions of eqs. (1) are the integral curves of the Hamiltonian vector field

$$\mathbf{X}_H := \frac{\partial H}{\partial p_i} \frac{\partial}{\partial q^i} - \frac{\partial H}{\partial q^i} \frac{\partial}{\partial p_i} \quad (3)$$

on $\mathcal{M}$. It is well known that the above ordinary Hamiltonian system preserves both the symplectic 2-form ω and the volume form $\frac{1}{n!} \omega^n$, where $n = \frac{1}{2} \dim \mathcal{M}$, and ω^k is the k -fold wedge product of ω .

It is well known that the ordinary Hamilton equations can be expressed as

$$-i_{\mathbf{X}_H} \omega = dH, \quad (4)$$

where i denotes the contraction. It is easy to show the symplectic structure-preserving and volume-preserving properties for these ordinary Hamiltonian systems. Namely,

$$\mathcal{L}_{\mathbf{X}_H} \omega = 0 \quad (5)$$

and

$$\mathcal{L}_{\mathbf{X}_H} (\omega^n) = n (\mathcal{L}_{\mathbf{X}_H} \omega) \wedge \omega^{n-1} = 0, \quad (6)$$

respectively.

However, there exist certain important mechanical systems that preserve the volume of the phase space only and cannot be transformed into the ordinary Hamiltonian systems. For example, the linear systems on $\mathbb{R}^n$

$$\ddot{q}^i = -k_{ij} q^j \quad (7)$$

¹ Repeated indices are always assumed to be summed over 1 to n unless otherwise stated. For convenience, symbols such like $\omega|_U$ are always abbreviated as ω .

with constant coefficients k_{ij} are such kind of mechanical systems. Obviously, eqs. (7) can be turned into the form:

$$\dot{q}^i = \frac{\partial H}{\partial p_i}, \quad \dot{p}_i = -\frac{\partial H}{\partial q^i} - a_{ij} q^j \quad (8)$$

with

$$H = \frac{1}{2} \delta^{ij} p_i p_j + \frac{1}{2} s_{ij} q^i q^j, \quad (9)$$

$$s_{ij} = \frac{1}{2} (k_{ij} + k_{ji}), \quad a_{ij} = \frac{1}{2} (k_{ij} - k_{ji}). \quad (10)$$

When one of a_{ij} is nonzero², the system (8) is not an ordinary Hamiltonian system on $\mathbb{R}^{2n}$. In fact, the corresponding vector field

$$\mathbf{X} = \mathbf{X}_H - a_{ij} q^j \frac{\partial}{\partial p_i}$$

of (8) is not even a symplectic vector field since the Lie derivative

$$\mathcal{L}_{\mathbf{X}}\omega = a_{ij} dq^i \wedge dq^j$$

does not vanish in this case. On the other hand, the linear system (8) always preserves the volume form of the phase space because $\mathcal{L}_{\mathbf{X}}(\omega^n) = 0$, as what can be easily verified.

In this letter, we present a general form of equations that generate a volume-preserving flow on a symplectic manifold $(\mathcal{M}, \omega)$. It is shown that every volume-preserving flow has some 2-forms acting the rôle of the Hamiltonian functions in the Hamiltonian mechanics and the ordinary Hamiltonian equations are included as a special case with a 2-form $\frac{1}{n-1} H \omega$ where H is the corresponding Hamiltonian.

² Here is an example showing that even a conservative system can be transformed to be a non-Hamiltonian system. Consider a system consisting of two 1-dimensional linearly coupled oscillators:

$$m_1 \ddot{q}^1 = -k(q^2 - q^1), \quad m_2 \ddot{q}^2 = -k(q^1 - q^2).$$

Obviously, such a system satisfies Newton's laws. Let $k_{11} = -k/m_1$, $k_{12} = k/m_1$, $k_{21} = k/m_2$ and $k_{22} = -k/m_2$. Then it is a system as described by eqs. (7), with the indices running from 1 to 2. When $m_1 \neq m_2$, we have a system satisfying $k_{12} \neq k_{21}$.

The main idea for presenting such general equations for volume-preserving systems is based on the linear isomorphism between the space $\mathcal{X}_H^n(\mathcal{M}, \omega)$ of certain volume-preserving vector fields and the space $B^{2n-1}(\mathcal{M})$ of exact $(2n-1)$ -forms on $\mathcal{M}$. This is, in fact, a direct generalization for the ordinary Hamiltonian system ($n = 1$). Namely, the linear isomorphism between the space $\mathcal{X}_H^{n=1}(\mathcal{M}, \omega)$ of certain symplectic structure-preserving vector fields and the space $B^1(\mathcal{M})$ of exact 1-forms on $\mathcal{M}$.

Briefly speaking, a volume-preserving system, described by a volume-preserving vector field with respect to which the Lie derivative of ω^n being zero, corresponds to a unique closed $(2n-1)$ -form on $\mathcal{M}$ and *vice versa*. Such a system that corresponds to an exact $(2n-1)$ -form can be determined then by a 2-form (may up to a closed 2-form), as if a Hamiltonian system can be determined by a function (the Hamiltonian) up to a constant function.

In this letter, the main results will be quoted and outlined without detailed proof. For the details and extended topics, we refer to our forthcoming papers.

For each integer $k = 1, \dots, n$ where $n = \frac{1}{2} \dim \mathcal{M} \geq 1$, we may define the set of the symplectic vector fields and that of the Hamiltonian vector fields as

$$\begin{aligned}\mathcal{X}_S^k(\mathcal{M}, \omega) &= \{ \mathbf{X} \in \mathcal{X}(\mathcal{M}) \mid \mathcal{L}_{\mathbf{X}}(\omega^k) = 0 \}, \\ \mathcal{X}_H^k(\mathcal{M}, \omega) &= \{ \mathbf{X} \in \mathcal{X}(\mathcal{M}) \mid -i_{\mathbf{X}}(\omega^k) \text{ is exact} \},\end{aligned}$$

respectively. Here and hereafter $\mathcal{X}(\mathcal{M})$ denotes the space of smooth vector fields on $\mathcal{M}$. Then it is easy to prove that

$$\begin{aligned}\mathcal{X}_S^1(\mathcal{M}, \omega) &\subseteq \dots \subseteq \mathcal{X}_S^k(\mathcal{M}, \omega) \subseteq \mathcal{X}_S^{k+1}(\mathcal{M}, \omega) \subseteq \dots \subseteq \mathcal{X}_S^n(\mathcal{M}, \omega), \\ \mathcal{X}_H^1(\mathcal{M}, \omega) &\subseteq \dots \subseteq \mathcal{X}_H^k(\mathcal{M}, \omega) \subseteq \mathcal{X}_H^{k+1}(\mathcal{M}, \omega) \subseteq \dots \subseteq \mathcal{X}_H^n(\mathcal{M}, \omega)\end{aligned}$$

and

$$\mathcal{X}_H^k(\mathcal{M}, \omega) \subseteq \mathcal{X}_S^k(\mathcal{M}, \omega).$$

In fact, $\mathcal{X}_S^k(\mathcal{M}, \omega)$ is a Lie algebra under the commutation bracket of vector fields, and $\mathcal{X}_H^k(\mathcal{M}, \omega)$ is an ideal of $\mathcal{X}_S^k(\mathcal{M}, \omega)$ because

$$[\mathcal{X}_S^k(\mathcal{M}, \omega), \mathcal{X}_H^k(\mathcal{M}, \omega)] \subseteq \mathcal{X}_H^k(\mathcal{M}, \omega).$$

Let us consider the quotient Lie algebra

$$H_{\text{EL}}^k(\mathcal{M}, \omega) := \mathcal{X}_S^k(\mathcal{M}, \omega) / \mathcal{X}_H^k(\mathcal{M}, \omega). \quad (11)$$

It can be called the k -th *Euler-Lagrange cohomology group*. It is called a group because its Lie algebra structure is trivial: For each $k = 1, \dots, n$, $H_{\text{EL}}^k(\mathcal{M}, \omega)$ is an abelian Lie algebra. For $n = 1$, $H_{\text{EL}}^1(\mathcal{M}, \omega)$ is the Euler-Lagrange cohomology first introduced in [2, 3]. And it can be proved that $H_{\text{EL}}^1(\mathcal{M}, \omega)$ is linearly isomorphic to the first de Rham cohomology group $H_{\text{dR}}^1(\mathcal{M})$.

For $k = n$, $\mathcal{X}_S^n(\mathcal{M}, \omega)$ is the Lie algebra of volume-preserving vector fields, including all the other Lie algebras mentioned in the above as its Lie subalgebras. Similarly to the case of $k = 1$, the n -th Euler-Lagrange cohomology group $H_{\text{EL}}^n(\mathcal{M}, \omega)$ is isomorphic to the $(2n - 1)$ -th de Rham cohomology group $H_{\text{dR}}^{2n-1}(\mathcal{M})$. To prove this, let us first introduce a lemma:

Lemma 1 *For each integer $k = 1, \dots, n$ ($n \geq 1$) and $x \in \mathcal{M}$, the linear map $\nu_k : T_x \mathcal{M} \longrightarrow \Lambda_{2k-1}(T_x^* \mathcal{M})$ defined by $\nu_k(\mathbf{X}) = -i_{\mathbf{X}}(\omega^k)$ is injective, namely, $i_{\mathbf{X}}(\omega^k) = 0$ iff $\mathbf{X} = 0$.*

Since $\dim T_x \mathcal{M} = \dim \Lambda_{2n-1}(T_x^* \mathcal{M})$, lemma 1 implies that $\nu_n : T_x \mathcal{M} \longrightarrow \Lambda_{2n-1}(T_x^* \mathcal{M})$ is a linear isomorphism, which induces a linear isomorphism $\nu_n : \mathcal{X}(\mathcal{M}) \longrightarrow \Omega^{2n-1}(\mathcal{M})$ defined by

$$\nu_n(\mathbf{X}) = -i_{\mathbf{X}}(\omega^n) = -n(i_{\mathbf{X}}\omega) \wedge \omega^{n-1}. \quad (12)$$

Hence we obtain

Theorem 1 *The linear map $\nu_n : \mathcal{X}(\mathcal{M}) \longrightarrow \Omega^{2n-1}(\mathcal{M})$ is an isomorphism. Under this isomorphism, $\mathcal{X}_S^n(\mathcal{M}, \omega)$ and $\mathcal{X}_H^n(\mathcal{M}, \omega)$ are isomorphic to $Z^{2n-1}(\mathcal{M})$ and $B^{2n-1}(\mathcal{M})$, respectively.*

In the above theorem, $Z^{2n-1}(\mathcal{M})$ is the space of closed $(2n - 1)$ -forms on $\mathcal{M}$ and $B^{2n-1}(\mathcal{M})$ is the space of exact $(2n - 1)$ -forms on $\mathcal{M}$. As a consequence, we obtain the following corollary:

Corollary 1 *The n -th Euler-Lagrange cohomology group $H_{\text{EL}}^n(\mathcal{M}, \omega)$ is linearly isomorphic to $H_{\text{dR}}^{2n-1}(\mathcal{M})$, the $(2n - 1)$ -th de Rham cohomology group.*

When $\mathcal{M}$ is closed (i.e., compact and without boundary), such as a closed Riemann surface, $H_{\text{EL}}^n(\mathcal{M}, \omega)$ is linearly isomorphic to the dual space of $H_{\text{EL}}^1(\mathcal{M})$, because $H_{\text{dR}}^k(\mathcal{M}) \cong (H_{\text{dR}}^{2n-k}(\mathcal{M}))^*$ for such a manifold (see, for example, [4]). If $\mathcal{M}$ is not compact, this relation cannot be assured.

When $n > 2$, there is another lemma:

Lemma 2 *If $n > 2$, then, for an arbitrary integer $k = 1, \dots, n-2$ and a point $x \in \mathcal{M}$, $\alpha \in \Lambda_2(T_x^* \mathcal{M})$ satisfies $\alpha \wedge \omega^k = 0$ iff $\alpha = 0$.*

Again this induces injective linear maps

$$\begin{aligned} \iota_k : \Omega^2(\mathcal{M}) &\longrightarrow \Omega^{2k+2}(\mathcal{M}) \\ \alpha &\longmapsto \alpha \wedge \omega^k \end{aligned}$$

for $k = 1, \dots, n-2$. Thus we can obtain that

$$\mathcal{X}_{\text{S}}^1(\mathcal{M}, \omega) = \mathcal{X}_{\text{S}}^2(\mathcal{M}, \omega) = \dots = \mathcal{X}_{\text{S}}^{n-1}(\mathcal{M}, \omega) \subseteq \mathcal{X}_{\text{S}}^n(\mathcal{M}, \omega).$$

When $k = n-2$, it can be easily verified from lemma 2 that

$$\begin{aligned} \iota = \iota_{n-2} : \Omega^2(\mathcal{M}) &\longrightarrow \Omega^{2n-2}(\mathcal{M}) \\ \alpha &\longmapsto \alpha \wedge \omega^{n-2} \end{aligned} \tag{13}$$

is a linear isomorphism. If $n = 2$, we use the convention that $\omega^0 = 1$, namely, $\iota = \text{id} : \alpha \longmapsto \alpha$. Then we can define a linear map ϕ making the following diagram commutative:

$$\begin{array}{ccc} \Omega^2(\mathcal{M}) & \xrightarrow{\iota} & \Omega^{2n-2}(\mathcal{M}) \\ \phi \downarrow & & \text{d} \downarrow \\ \mathcal{X}_{\text{H}}^n(\mathcal{M}, \omega) & \xrightarrow{\nu_n} & B^{2n-1}(\mathcal{M}) \end{array} \tag{14}$$

Equivalently, given a 2-form

$$\alpha = \frac{1}{2} Q_{ij} dq^i \wedge dq^j + A_j^i dp_i \wedge dq^j + \frac{1}{2} P^{ij} dp_i \wedge dp_j \tag{15}$$

where Q_{ij} and P^{ij} satisfy

$$Q_{ji} = -Q_{ij}, \quad P^{ji} = -P^{ij}, \tag{16}$$

the vector field

$$\phi(\alpha) = (\nu_n^{-1} \circ d \circ \iota)(\alpha) = \nu_n^{-1}(d\alpha \wedge \omega^{n-2})$$

is in $\mathcal{X}_H^n(\mathcal{M}, \omega)$.

For convenience, we set

$$\mathbf{X} = n(n-1) \phi(\alpha) = n(n-1) \nu_n^{-1}(d\alpha \wedge \omega^{n-2}),$$

namely, $i_{\mathbf{X}}(\omega^n) = -n(n-1) (d\alpha) \wedge \omega^{n-2}$. It is easy to obtain that

$$\mathbf{X} = \left(\frac{\partial P^{ij}}{\partial q^j} + \frac{\partial A_j^j}{\partial p_i} - \frac{\partial A_j^i}{\partial p_j} \right) \frac{\partial}{\partial q^i} + \left(\frac{\partial Q_{ij}}{\partial p_j} - \frac{\partial A_j^j}{\partial q^i} + \frac{\partial A_i^j}{\partial q^j} \right) \frac{\partial}{\partial p_i}. \quad (17)$$

In fact,

$$\begin{aligned} d\iota(\alpha) &= d\alpha \wedge \omega^{n-2} \\ &= \frac{1}{2} \frac{\partial Q_{ij}}{\partial q^k} dq^i \wedge dq^j \wedge dq^k \wedge \omega^{n-2} + \frac{1}{2} \frac{P^{ij}}{\partial p_k} dp_i \wedge dp_j \wedge dp_k \wedge \omega^{n-2} \\ &\quad + \left(\frac{1}{2} \frac{\partial Q_{jk}}{\partial p_i} + \frac{\partial A_j^i}{\partial q^k} \right) dp_i \wedge dq^j \wedge dq^k \wedge \omega^{n-2} \\ &\quad + \left(\frac{1}{2} \frac{\partial P^{ij}}{\partial q^k} - \frac{\partial A_k^i}{\partial p_j} \right) dp_i \wedge dp_j \wedge dq^k \wedge \omega^{n-2} \\ &= \left(\frac{1}{2} \frac{\partial Q_{jk}}{\partial p_i} + \frac{\partial A_j^i}{\partial q^k} \right) dp_i \wedge dq^j \wedge dq^k \wedge \omega^{n-2} \\ &\quad + \left(\frac{1}{2} \frac{\partial P^{ij}}{\partial q^k} - \frac{\partial A_k^i}{\partial p_j} \right) dp_i \wedge dp_j \wedge dq^k \wedge \omega^{n-2}. \end{aligned}$$

By virtue of the following two equations

$$dp_i \wedge dp_j \wedge dq^k \wedge \omega^{n-2} = \frac{\delta_j^k}{n-1} dp_i \wedge \omega^{n-1} - \frac{\delta_i^k}{n-1} dp_j \wedge \omega^{n-1}, \quad (18)$$

$$dp_i \wedge dq^j \wedge dq^k \wedge \omega^{n-2} = \frac{\delta_i^j}{n-1} dq^k \wedge \omega^{n-1} - \frac{\delta_i^k}{n-1} dq^j \wedge \omega^{n-1}, \quad (19)$$

we can write $d\iota(\alpha)$ as

$$d\iota(\alpha) = \frac{1}{n-1} \left(\frac{\partial A_j^j}{\partial q^i} - \frac{\partial A_i^j}{\partial q^j} - \frac{\partial Q_{ij}}{\partial p_j} \right) dq^i \wedge \omega^{n-1}$$

$$\begin{aligned}
& + \frac{1}{n-1} \left(\frac{\partial P^{ij}}{\partial q^j} + \frac{\partial A_j^j}{\partial p_i} - \frac{\partial A_j^i}{\partial p_j} \right) dp_i \wedge \omega^{n-1} \\
& = \frac{1}{n(n-1)} \left(\frac{\partial A_j^j}{\partial q^i} - \frac{\partial A_i^j}{\partial q^j} - \frac{\partial Q_{ij}}{\partial p_j} \right) i_{\frac{\partial}{\partial p_i}} \omega^n \\
& \quad - \frac{1}{n(n-1)} \left(\frac{\partial P^{ij}}{\partial q^j} + \frac{\partial A_j^j}{\partial p_i} - \frac{\partial A_j^i}{\partial p_j} \right) i_{\frac{\partial}{\partial q^i}} \omega^n.
\end{aligned}$$

Comparing it with

$$d\iota(\alpha) = \frac{1}{n(n-1)} \nu_n(\mathbf{X}) = -\frac{1}{n(n-1)} i_{\mathbf{X}}(\omega^n), \quad (20)$$

we can obtain the expression of $\mathbf{X}$, as shown in eq. (17).

Since both ι and ν_n are linear isomorphisms, we can see from the commutative diagram (14) that for each 2-form α on $\mathcal{M}$ as in eq. (15) the vector field $\mathbf{X}$ in eq. (17) belongs to $\mathcal{X}_{\text{H}}^n(\mathcal{M}, \omega)$ and for each $\mathbf{X} \in \mathcal{X}_{\text{H}}^n(\mathcal{M}, \omega)$ there exists the 2-form α on $\mathcal{M}$ satisfying eqs. (17). But there may be several 2-forms that are mapped to the same vector field $\mathbf{X}$. For example, the vector field $\mathbf{X}$ in (17) is invariant under the transformation

$$\alpha \longmapsto \alpha + \theta \quad (21)$$

where θ is a closed 2-form.

Note that for the vector field $\mathbf{X} \in \mathcal{X}_{\text{H}}^n(\mathcal{M}, \omega)$, the 2-form α is a globally defined differential form on $\mathcal{M}$. If $\mathbf{X} \in \mathcal{X}_{\text{S}}^n(\mathcal{M}, \omega)$, such a 2-form cannot be found if $H_{\text{dR}}^{2n-1}(\mathcal{M})$ is nontrivial. In this case, α can be still found as a locally defined 2-form, according to Poincaré's lemma. Then the relation between $\mathbf{X}$ and the locally defined 2-form α , eq. (17), is valid only on a certain open subset of $\mathcal{M}$, and, on the intersection of two such open subsets, the corresponding 2-forms are not identical. That is, the transformation relation of Q_{ij} , A_j^i as well as P^{ij} is not that of tensors on $\mathcal{M}$. Instead, a transformation such like (21) or more complicated should be applied.

No matter whether $\mathbf{X}$ belongs to $\mathcal{X}_{\text{H}}^n(\mathcal{M}, \omega)$ or $\mathcal{X}_{\text{S}}^n(\mathcal{M}, \omega)$, namely, whether the 2-form α is globally or locally defined, the flow of $\mathbf{X}$ can be always obtained provided that the general solution of the following equations can be solved:

$$\dot{q}^i = \frac{\partial P^{ij}}{\partial q^j} + \frac{\partial A_j^j}{\partial p_i} - \frac{\partial A_j^i}{\partial p_j},$$

$$\dot{p}_i = \frac{\partial Q_{ij}}{\partial p_j} - \frac{\partial A_j^i}{\partial q^i} + \frac{\partial A_i^j}{\partial q^j}. \quad (22)$$

This is just the general form of the equations of a volume-preserving mechanical system on a symplectic manifold $(\mathcal{M}, \omega)$.

For a function H on $\mathcal{M}$, let the 2-form be

$$\alpha = \frac{1}{n-1} H \omega, \quad (23)$$

then eqs. (22) turn to be the ordinary Hamilton equations (1). Thus the ordinary Hamilton equations have been included as a special case, as what is expected.

Now suppose the symplectic manifold is $\mathbb{R}^{2n}$ with the standard symplectic form $\omega = dp_i \wedge dq^i$. When $Q_{ij} = -a_{ij} p_k q^k$, $A_j^i = \frac{1}{n-1} H \delta_j^i$ and $P^{ij} = 0$ with the constants a_{ij} and the function H as shown in eqs. (9) and (10), namely,

$$\alpha = \frac{1}{n-1} H \omega - \frac{1}{2} p_k q^k a_{ij} dq^i \wedge dq^j, \quad (24)$$

then the equations (22) will turn into eqs. (8). Therefore the 2-form α in eq. (24) is one of general forms corresponding to a linear system, but it is not the unique.

It is interesting that when $H = 0$ in the above equation, all the coordinates q^i are the first integrals of the linear system. Then all the canonical momenta $p_i = a_{ij} q^j t + p_{i0}$ where p_{i0} are constants. This can generalize to an arbitrary symplectic manifold for the equations (22), even though the system is no longer a linear system: If, on a Darboux coordinate neighborhood $(U; q, p)$ in a symplectic manifold $(\mathcal{M}, \omega)$, the 2-form α satisfies

$$i_{\frac{\partial}{\partial p_i}} \alpha = A_j^i dq^j + P^{ij} dp_j = 0 \quad (25)$$

for each $i = 1, \dots, n$, then $A_j^i = 0$ and $P^{ij} = 0$. Hence, on that coordinate neighborhood U , all the q^i are constant.

If we define a function $\text{tr } \alpha$ as

$$\alpha \wedge \omega^{n-1} = \frac{\text{tr } \alpha}{n} \omega^n \quad (26)$$

for each 2-form α , then using the formula

$$dp_i \wedge dq^j \wedge \omega^{n-1} = \frac{\delta_i^j}{n} \omega^n,$$

we obtain that

$$\mathrm{tr} \alpha = A_i^i. \quad (27)$$

The above expression is obviously independent of the choice of the Darboux coordinates. Let $\mathbf{X}_{\mathrm{tr} \alpha}$ be the Hamiltonian vector field corresponding to the function $\mathrm{tr} \alpha$. Eq. (17) indicates that

$$\mathbf{X} = \mathbf{X}_{\mathrm{tr} \alpha} + \mathbf{X}' \quad (28)$$

where $\mathbf{X}'$ is the extra part on the right hand side of eq. (17), corresponding to the 2-form

$$\alpha - \frac{\mathrm{tr} \alpha}{n-1} \omega. \quad (29)$$

If $f(q, p)$ is a function on $\mathcal{M}$, then the derivative $\dot{f} = \frac{d}{dt}f(q(t), p(t))$ along any one of the integral curves of eqs. (22) satisfies the equation

$$\dot{f} \omega^n = n(n-1) d\alpha \wedge df \wedge \omega^{n-2}. \quad (30)$$

In fact, $\dot{f}(t) = (\mathcal{L}_{\mathbf{X}}f)(q(t), p(t))$. And, since $\mathbf{X}$ is volume-preserving,

$$(\mathcal{L}_{\mathbf{X}}f) \omega^n = \mathcal{L}_{\mathbf{X}}(f \omega^n) = di_{\mathbf{X}}(f \omega^n) + i_{\mathbf{X}}d(f \omega^n) = d(f i_{\mathbf{X}}\omega^n).$$

Then according to eq. (20),

$$\begin{aligned} (\mathcal{L}_{\mathbf{X}}f) \omega^n &= -n(n-1) d(f d\iota(\alpha)) = -n(n-1) df \wedge d(\alpha \wedge \omega^{n-2}) \\ &= -n(n-1) df \wedge d\alpha \wedge \omega^{n-2} = n(n-1) d\alpha \wedge df \wedge \omega^{n-2}. \end{aligned}$$

Thus eq. (30) has been proved.

Especially, when $\alpha = \frac{H}{n-1} \omega$, the system (22) turns out to be the usual Hamiltonian system, as we have mentioned. For such a Hamiltonian system, on the one hand, we can use eq. (30) to obtain

$$\dot{f} \omega^n = n d(H \omega) \wedge df \wedge \omega^{n-2} = n dH \wedge df \wedge \omega^{n-1} = \mathrm{tr} (dH \wedge df) \omega^n,$$

namely,

$$\dot{f} = \mathrm{tr} (dH \wedge df). \quad (31)$$

On the other hand, $\dot{f}$ can be expressed in terms of the Poisson bracket:

$$\dot{f} = \{f, H\} := \mathbf{X}_H f = \frac{\partial f}{\partial q^i} \frac{\partial H}{\partial p_i} - \frac{\partial f}{\partial p_i} \frac{\partial H}{\partial q^i}.$$

So we obtain the relation between the Poisson bracket and the trace of 2-forms:

$$\{f, H\} = -\text{tr}(\text{d}f \wedge \text{d}H). \quad (32)$$

In order to develop a volume-preserving algorithm, Feng and Shang in [5] presented a lemma as follows: The volume-preserving vector field $X = (X^1, \dots, X^n)^T$ on $\mathbb{R}^n$ can always be expressed by an antisymmetric tensor a^{ij} on $\mathbb{R}^n$ as³

$$X^i = \frac{\partial a^{ij}}{\partial x^j}, \quad (33)$$

where x^i are the standard coordinates on $\mathbb{R}^n$. In fact, this lemma is not difficult to understand: A vector field on $\mathbb{R}^n$, known as a Euclidean space or a Riemannian manifold, is volume-preserving if and only if it is divergence-free. Using the Hodge theory, the divergence of the vector field X reads $\text{div} X = -\delta \tilde{X} = * \text{d} * \tilde{X}$, where δ is the codifferential operator and $\tilde{X} = \delta_{ij} X^i \text{d}x^j$ is the 1-form, or the covariant vector field, obtained from X by using Einstein's method of lowering indices. Hence that X is volume-preserving, namely, $\text{div} X = * \text{d} * \tilde{X} = 0$, implies $*\tilde{X}$ is a closed $(n-1)$ -form. Then, according to Poincaré's lemma, there is a 2-form α , say, such that $*\tilde{X} = \text{d} * \alpha$. If α is assumed to be $\frac{1}{2} a_{ij} \text{d}x^i \wedge \text{d}x^j$ with $a_{ij} = -a_{ji} = \delta_{ik} \delta_{jl} a^{kl} = a^{ij}$, we can obtain that $\tilde{X} = (-1)^{n-1} * \text{d} * \alpha = \delta_{ij} \frac{\partial a^{jk}}{\partial x^k} \text{d}x^i$. Then the vector field X satisfies eq. (33).

It is worthy to see that the formula (33) is quite similar to eq. (17), only with some slight differences: 1. This formula is a statement on the Euclidean space $\mathbb{R}^n$ with an arbitrary dimension n while eq. (17) is for a symplectic manifold, of $2n$ -dimensional. 2. It can be generalized to an n -dimensional Riemannian or pseudo-Riemannian manifold provided that the $(n-1)$ -th de Rham cohomology group is trivial, with the ordinary derivatives being replaced by the covariant derivatives. As for eq. (17), when $H_{\text{dR}}^{2n-1}(\mathcal{M}) \neq 0$, not every volume-preserving vector field satisfies it. This is similar to eq. (33). But a covariant derivative is not necessary in eq. (17). 3. When the symplectic manifold $\mathcal{M} = \mathbb{R}^{2n}$, we can set

$$(a_{ij})_{2n \times 2n} = \begin{pmatrix} 0 & I \\ -I & 0 \end{pmatrix} \begin{pmatrix} Q & -A^T \\ A & P \end{pmatrix} \begin{pmatrix} 0 & -I \\ I & 0 \end{pmatrix} = \begin{pmatrix} P & -A \\ A^T & Q \end{pmatrix}$$

³ In the quotation of this lemma, we have adopted the notations so as to accommodate ours in this letter.

where $Q = (Q_{ij})_{i=1, j=1}^n$, $P = (P^{ij})_{i=1, j=1}^n$, $A = (A_j^i)_{i=1, j=1}^n$ and I is the $n \times n$ unit matrix. Accordingly, we set $(x^1, \dots, x^n, x^{n+1}, \dots, x^{2n}) = (q^1, \dots, q^n, p_1, \dots, p_n)$. Then eq. (33) reads

$$X = \left(\frac{\partial P^{ij}}{\partial q^j} - \frac{\partial A_j^i}{\partial p_j}, \frac{\partial A_i^j}{\partial q^j} + \frac{\partial Q_{ij}}{\partial p_j} \right)^T, \quad (34)$$

with the index i running from 1 to n . Comparing it with eq. (17), it seems that a condition

$$\text{tr } \alpha = 0 \quad (35)$$

could be imposed on the 2-form α in eq. (17).

Finally, it should be mentioned that the volume-preserving systems play an important rôle in physics, especially in statistical physics. However, this has not been noticed widely yet. In this letter we have not explained why the volume-preserving systems are so important in statistical physics, but left it as a topic for the forthcoming papers.

Acknowledgement

We would like to thank Professors J. Z. Pan and Z. J. Shang for valuable discussions. This work was supported in part by the National Natural Science Foundation of China (grant Nos. 90103004, 10175070) and the National Key Project for Basic Research of China (G1998030601).

References

- [1] V. I. Arnold, *Mathematical Methods of Classical Mechanics*, Springer-Verlag, New-York, 1989.
- [2] H. Y. Guo, Y. Q. Li and K. Wu, “On Symplectic and Multisymplectic Structures and Their Discrete Version in Lagrangian Formalism”, *Comm. Theor. Phys.* (Beijing) **35** (2001) 703-710.
- [3] H. Y. Guo, Y. Q. Li, K. Wu and S. K. Wang, “Difference Discrete Variational Principle, Euler-Lagrange Cohomology and Symplectic, Multisymplectic structures I: Difference Discrete Variational Principle”, *Comm. Theor. Phys.* (Beijing), **37** (2002) 1-10; “II: Euler-Lagrange

Cohomology”, *ibid*, 129-138; “III: Application to Symplectic and Multisymplectic Algorithms”, *ibid*, 257-264. See, also, the eprint [hep-th/0106001](#).

- [4] F. W. Warner, *Foundations of Differentiable Manifolds and Lie Groups*, Springer-Verlag, New York, 1983.
- [5] Feng Kang and Shang Zai-jiu, “Volume-preserving algorithms for source-free dynamical systems”, *Numer. Math.*, Vol. **71** (1995) 451-463.